



King Saud University

Saudi Journal of Biological Sciences

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Ecology of the Sand Boa, *Eryx jayakari* in Riyadh Region of Saudi Arabia



Mohammed K. Al-Sadoon *, Fahed S. Al-Otaibi

Zoology Department, College of Science, King Saud University, PO Box 2455, Riyadh 11451, Saudi Arabia

Received 7 April 2014; revised 22 April 2014; accepted 30 April 2014

Available online 9 May 2014

KEYWORDS

Reptiles;
 Arabian Sand Boa;
 Feeding habitat;
 Sexual dimorphism

Abstract The ecology, feeding habits and sexual dimorphism in Arabian Sand Boa, *Eryx jayakari* from the Central region of Saudi Arabia, were studied. In this study the *E. jayakari* was recorded for the first time from several sites. Significant differences were noted in total length of body and tail, and body diameter of male and female. The females were of larger size. The mean number of the dorsal body scales, ventrals and subcaudal for both sexes were not significantly different. The mean number of the dorsal body scales, ventral body scales and subcaudal scales for the females was 43, 169 and 18 scales which were not significantly different from respectively ones in males 42, 168 and 18 scales. Frequent prey consumed were lizards (50%), rodents (25%) and arthropods (12.5%).

© 2014 Production and hosting by Elsevier B.V. on behalf of King Saud University.

1. Introduction

Saudi Arabia lies between Africa and Asia, occupying about three fourth of the Arabian Peninsula. Saudi Arabia is hot (50 °C) in summer and cold (20 °C) in winter. Saudi Arabia lies in the tropical and sub-tropical region, it has – mountains, vast sandy deserts, valleys and forests. Despite hostile environments, a considerable number of species of reptiles occur here.

Few ecological studies have been done on snakes and lizards of different regions of Saudi Arabia *viz.*, Southern Hijaz (Parker, 1933, 1938), Eastern Arabia and Northeastern

Arabia (Mandaville, 1965, 1967), Central Arabia (Schmidt, 1941; Al-Wailly and Al-Uthman, 1971), and Riyadh (Hussein, 1966). Farag and Banaja (1980) have identified four anuran species, 28 species of lizards and 15 species of snakes, mapping their distribution in the Western region of Saudi Arabia.

Of the 51 species of snakes, nine are sea snakes, 42 are terrestrial, of which 9 are venomous (Mandaville, 1965; Gasperetti, 1974, 1988; Leviton, 1977a,b; Al-Sadoon, 1989). Reptiles have low ability to withstand changes in habitat; their dispersal is particularly dependent on environmental conditions. Few species have been adopted to peculiarities of habitats in the Sahara desert (Abuzinada et al., 2004). Snakes play important role in the ecosystem, they control rodent and insect populations in check (Masood, 2012).

The Arabian Sand Boa, *Eryx jayakari* is a small harmless snake, only boid found in the Arabian Peninsula. It is nocturnal, lives mostly under the desert sand, eyes are positioned on the top of head, rather than on sides (Fig. 1). It allows the snake to see while body remains buried in the sand. *E. jayakari*

* Corresponding author. Tel.: +966 505421754; fax: +966 114678514.

E-mail address: msadoon@ksu.edu.sa (M.K. Al-Sadoon).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Figure 1 The Sand Boa, *Eryx jayakari*.

is carnivores, it preys mainly on small rodents, reptiles and arthropods (Al-Johany and Al-Sadoon, 1996; Hellyer and Aspinall, 2005).

Unlike most boas, which give birth to live young, the Arabian Sand Boa is one of the three boa species that lay eggs (Bartlett and Bartlett, 2005). Usually a clutch of seven eggs is laid, which take around two months to hatch (Hellyer and Aspinall, 2005).

In body coloration it appears similar to the more familiar East African Sand Boa, *Eryx colubrinus*, with a black or brown pattern on an orange/yellow background. However, the pattern is finer and more irregular than that of *E. colubrinus* and the darker blotches may form irregular bands across the body of the snake. In some parts of its range the Arabian Sand Boas are darker and almost pattern less, a condition similar to that seen on some populations of *Eryx miliaris*, *Eryx conicus* and *E. colubrinus*. The present work is a detailed study of ecology and biology of *E. jayakari* in Riyadh Region.

2. Materials and methods

The field study was carried out in different parts of the sandy regions (Nafud Qunayfidhan, Nafud al Sirr, Nafud Dahna, Nafud Al Mazhur, Saad Park and Al-Thumamah) at Riyadh, Saudi Arabia; in a diameter 200 km in the four directions, for one year on monthly basis, from March 2010 to February 2011. During the survey the snake tracks were followed to reach them in their burrows. The data are mapped using the GPS. Ecological field studies including distribution, behavioral, morphological characters and the mode of nutrition were carried out.

Seventy animals were collected for laboratory studies. The animals were killed by freezing at -2°C for 24 h. The parameters recorded were: the body weight, snout-vent length, vent-tail length, total length and diameter according to Schoener (1968), number of dorsal scales at mid-body and ventral scales. The digestive tract was removed and preserved in 70% ethanol for investigation on the food contents.

3. Results and discussion

E. jayakari (Fig. 1) was found in several sites in Riyadh viz., Thumamah, the eastern and northern Adhirah, Saad and

Wasia (East), Nfod Alzoillat (South), Nfod Hammad area (North), near the Al Zulfi area, El Muzahmiyya area (west), and southern Nfod Guenivzh.

Table 1 shows significant difference between both the males and females in terms of total length, snout-vent length and body diameter with the females attaining larger size ($P < 0.05$). The mean number of the dorsal body scales, ventral body scales and subcaudal scales for the females was 43, 169 and 18 scales which were not significantly different from respective ones in males 42, 168 and 18 scales (Table 1).

The result of stomach contents indicates that *E. jayakari* is carnivorous and feeds mainly on: lizards (*Acanthodactylus schmidti*, *Bumopus tuberculatus* and *Stenodactylus slevinii*) which form 50% of the stomach contents, rodents (*Gerbillus cheesmani* and *Gerbillus nanus*) in snakes with larger size and formed 25% of the stomach contents; and Arthropods (beetles) constituting 12.5% of the content. The remaining 12.5% of the stomach contents was completely digested and could not be identified (Fig. 2).

In this study the *E. jayakari* was recorded for the first time from several sites. Our results reveal that *E. jayakari* is a nocturnal forager, feeds on a variety of animals as concluded by Mattison (1995) and Gasperetti (1988).

The female *E. jayakari* are larger than the males (Fitch, 1960, 1975; Wilbur, 1975; Vogt, 1980), exhibit sexual dimorphism in size and growth rate. Sex differences in body size are widespread in snakes, mostly females are larger than males (Shine, 1994).

Larger animals tend to feed upon larger prey, and hence leading to decreased interspecific and intraspecific competition (Best and Gennaro, 1984).

Table 1 Body measurements and the number of scales of male and females in *E. jayakari*. (M \pm SE).

Measurements	Male (n = 45)	Female (n = 25)
<i>Body</i>		
Snout-vent length	318 \pm 6.17	372 \pm 11.7
Vent-tail length	20.4 \pm 0.60	20.2 \pm 0.60
Total length	388 \pm 6.85	390 \pm 12
Diameter	10.5 \pm 0.22	13 \pm 0.29
<i>Scales</i>		
Dorsal scales	42 \pm 0.20	43 \pm 0.19
Ventral scales	168 \pm 0.19	169 \pm 0.22
Sub-caudal scales	18.6 \pm 0.14	18 \pm 0.19




Figure 2 Stomach contents of *E. jayakari*.

Although *E. jayakari* is nocturnal, the existence of diurnally active lizards in the stomach may be due to snakes forage at night for inactive prey. Shine (1982) has also reported nocturnal foraging of inactive skinks as the feeding “strategy” of small Australian elapid snakes of the genera *Cacophis*, *Furina*, *Glyphodon*, *Neelaps*, *Unechis* and some species within *Denisonia* and *Simoselaps*.

Acknowledgment

The authors would like to express their sincere appreciation to the Deanship of Scientific Research at the King Saud University, Riyadh, Saudi Arabia for funding this Research Group project no. RGP-VPP-289.

References

- Abuzinada, A.H., Robinson, E.R., Nader, I.A., Al-Wetaid, Y.I., 2004. First Saudi Arabian National Report on the convention on Biological diversity. The National Commission for Wildlife Conservation and Development, Riyadh, Saudi Arabia, pp. 27–28.
- Al-Johany, A.M., Al-Sadoon, M.K., 1996. Selected body temperature and metabolic rate-temperature curves of three species of desert snakes. *J. Arid Environ.* 34, 363–370.
- Al-Sadoon, M.K., 1989. Survey of the reptilian fauna of the Kingdom of Saudi Arabia. I. The snake fauna of the Central Region. *J. King Saud Univ. 1A Sci.* 1, 53–69.
- Al-Wailly, A.J., Al-Uthman, H.S., 1971. Some lizards from Central Saudi Arabia. *Bull. Iraq Nat. Hist. Mus.* 5, 39–42.
- Bartlett, R.D., Bartlett, P.P., 2005. *Rosy, Rubber, and Sand Boas: Facts and Advice on Care and Breeding*. Barron's Educational Series, New York.
- Best, T.L., Gennaro, A.L., 1984. Feeding ecology of the lizard, *Uta stansburiana*, in southwestern Mexico. *J. Herpetol.* 18, 291–301.
- Farag, A.A., Banaja, A.A., 1980. Amphibians and reptiles from the western region of Saudi Arabia. *Bull. Sci. K. A. U.* 4, 5–29.
- Fitch, H.S., 1960. Autecology of the copperhead. *Univ. Kansas Publ. Mus. Natur. Hist.* 13, 85–288.
- Fitch, H.S., 1975. A demographic study of the ringneck snake (*Diadophis punctatus*) in Kansas. *Univ. Kansas Misc. Pub.* 62, 1–53.
- Gasperetti, J., 1974. A preliminary sketch of the snakes of the Arabian Peninsula. *J. Saudi Nat. Hist. Soc.* 12, IV + 72.
- Gasperetti, J., 1988. The Snakes of Arabia. *Fauna Saudi Arabia* 9, 169–450.
- Hellyer, P., Aspinall, S. (Eds.), 2005. *The Emirates. A Natural History*. Trident Press Ltd, Mayfair, London, UK, p. 312.
- Hussein, M.F., 1966. On the reptiles of Riyadh and its surroundings in Saudi Arabia. 5th Arab Sci. Congr. Baghdad., 479–485.
- Leviton, A.E., 1977a. A new lytorhynchid snake. *J. Saudi Nat. Hist. Soc.* 19, 16–25.
- Leviton, A.E., 1977b. On the status of *Natrix dubbiosi scorececi*. *J. Saudi Nat. Hist. Soc.* 19, 26–29.
- Mandaville, J.P., 1965. Plants eaten by *Uromastix microlepis* Blandford and other notes on these lizards in Eastern Arabia. *J. Bombay Nat. Hist. Soc.* 61 (10), 161–163.
- Mandaville, J.P., 1967. The hooded *Malpolon moileensis* reuss and notes on other snakes of North-Eastern Arabia. *J. Bombay Nat. Hist. Soc.* 64 (1), 115–117.
- Masood, F.M., 2012. Ecological distribution of snakes' fauna of Jazan region of Saudi Arabia. *Egypt. Acad. J. Biolog. Sci.* 4 (1), 183–197.
- Mattison, C., 1995. *The Encyclopedia Of Snakes*. Checkmark Books, New York, U.S.A., p. 265.
- Parker, H.W., 1933. Reptiles and amphibians. In: Philby, H.St.J.B. (Ed.), *The Empty Quarter*. Constable, London, pp. 97–398.
- Parker, H.W., 1938. Reptiles and amphibians from Southern Hejaz. *Ann. Mag. Nat. Hist.* 1 (5), 481–492 (Set. 11).
- Schmidt, K.P., 1941. Reptiles and Amphibia from Central Arabia. *Field Mus. Nat. Hist. Zool. Ser.* 24 (16), 161–165.
- Schoener, T.W., 1968. The Anolis lizards of Bimini: resource partitioning in a complex fauna. *Ecology* 49, 704–726.
- Shine, R., 1982. Ecology of the Australian elapid snake *Echiopsis curta*. *J. Herpetol.* 16, 388–393.
- Shine, R., 1994. Sexual size dimorphism in snakes revisited. *Copeia*, 326–346.
- Vogt, R.C., 1980. Natural history of the map turtles *Graptemys pseudogeographica* and *G. ouachitensis* in Wisconsin. *Tulane Stud. Zool. Bot.* 22, 17–48.
- Wilbur, H.M., 1975. The evolutionary and mathematical demography of the turtle *Chrysemys picta*. *Ecology* 56, 64–77.